

## OPINION

# The heat-shock response in higher plants: a biochemical model

J. J. BURKE & K. A. ORZECH USDA-ARS Plant Stress and Water Conservation Research Unit, Cropping Systems Research Laboratory, Box 215, Route 3, Lubbock, TX 79401, U.S.A.

Received 27 November 1987; accepted for publication 23 February 1988

**Abstract.** A compilation of existing data on higher plant responses to heat-shock temperatures has been utilized to produce a biochemically based model of integrated cellular responses to elevated temperatures. This model describes a potential mechanism for the triggering of several biochemical responses to a thermally induced leakage of extracellular or vacuolar ions into the cytoplasm. It seems possible that many of the observed heat-shock responses are involved in the protection of (a) enzymes from inactivation and (b) nucleic acids from cleavage induced by the presence of elevated levels of specific metals.

**Key-words:** heat shock; model; metals.

Living organisms have developed several endogenous protection systems which provide thermal tolerance. One of these protection systems involves an acquired heat resistance mechanism which is associated with the synthesis and accumulation of specific proteins (HSP). Universally these proteins have been identified by following the incorporation of labelled amino acids into proteins during exposure to elevated temperatures. The heat-shock response has been reviewed both at the protein and molecular levels in numerous articles, reviews and books (Baszczynski, Walden & Atkinson, 1985; Craig, 1986; Key *et al.*, 1982, 1983, 1985; Lindquist, 1986; Nover, 1984).

Early reports of heat-shock proteins in higher plants demonstrated HSP in tobacco and soybean cells grown in solution culture (Barnett *et al.*, 1980), and in soybean seedling tissue (Key, Lin & Chen, 1981). The HSP of soybean were demonstrated by the appearance of 10 new bands on one-dimensional SDS gels; with a more complex pattern on two-dimensional gels (Key *et al.*, 1981). When the tissue was returned to 28 °C after 4 h at 40 °C, there was a progressive decline in the synthesis of HSP and a reappearance of the normal pattern of protein synthesis by 3 to 4 h. To date, the incorporation of radioactive precursors into plant heat-shock proteins

has been identified in peas (Hadwiger & Wagoner, 1983; Mansfield & Key, 1987), tomato (Nover & Scharf, 1984; Nover, Scharf & Neumann, 1983), carrot (Pitto *et al.*, 1983), tobacco (Meyer & Chartier, 1983), mung bean (Chen, Kamisaka & Masuda, 1986), barley (Belanger, Brodl & Ho, 1986; Mansfield & Key, 1987), *Tradescantia* (Xiao & Mascarenhas, 1985), *Gladiolus* cormels (Ginzburg & Salomon, 1986), *Lilium longiflorum* (Hong-Qi, Croes & Linskens, 1984), soybean (Key *et al.*, 1981; Mansfield & Key, 1987), corn (Baszczynski, Walden & Atkinson, 1983; Bewley, Larson & Papp, 1983; Cooper & Ho, 1983, 1984; Mansfield & Key, 1987), cotton (Burke *et al.*, 1985), wheat (Key *et al.*, 1983; Mansfield & Key, 1987), millet (Key *et al.*, 1983; Mansfield & Key, 1987), sunflowers (Schoffl & Baumann, 1985), sorghum (Ougham & Stoddart, 1986), rice and *Panicum miliaceum* (Mansfield & Key, 1987). The optimal induction temperature for the heat-shock response varies between species, but generally occurs from 10 to 15 °C above the temperature empirically determined for optimal plant growth.

Several inducers of the synthesis of all or specific heat-shock proteins other than elevated temperatures have been identified in both plants and animals. These include developmental controls (Bienz, 1984), metals (Czarnecka *et al.*, 1984), water stress (Heikkila *et al.*, 1984), sulphhydryl reagents, calcium ionophores, steroid hormones, chelating agents, pyridoxine, methylene blue, glucosamine, deoxyglucose, and a variety of DNA and RNA viruses (Nover, 1984). Variability exists in the pattern of mRNA and proteins synthesized in response to the magnitude of the temperature shifts (Craig, 1986), chemical inducers (Lindquist, 1986), the degree to which 'normal' protein synthesis is inhibited (Key *et al.*, 1983), and the length of time that maximal HSP synthesis occurs (Baszczynski *et al.*, 1985). The reported cellular localizations of specific HSP in the nucleus (Vincent & Tanguay, 1979), mitochondria (Sinibaldi & Turpen, 1985; Cooper & Ho, 1987), chloroplast (Vierling *et al.*, 1986), endoplasmic reticulum (Baszczynski *et al.*, 1983; Cooper & Ho, 1987), and plasma membrane (Lim *et al.*, 1984; Cooper & Ho, 1987) can also vary depending upon

Correspondence: John J. Burke, USDA-ARS Plant Stress and Water Conservation Research Unit, Cropping Systems Research Laboratory, Box 215, Route 3, Lubbock, TX 79401, U.S.A.

the method of heat-shock protein induction (Lindquist, 1986; Craig, 1986).

One thread common to all of the heat-shock reviews is the inability of investigations to add insight into the question of heat-shock protein function. This paper presents a hypothesis of cellular responses to thermal stress based upon the existing literature, and suggests that some of the proteins associated with the heat-shock response may be associated with the protection of cellular processes from metal inhibition or toxicity. This manuscript is not intended to extensively review the existing literature, but will provide an overview based upon several aspects of the heat shock response in plants. Figure 1 presents a working model of potential cellular responses to high temperature stress.

In this model, elevation of cellular temperature would result in a decreased efficiency of membrane transport systems in the removal of metals and salts from the cytoplasm. This decreased efficiency could be related to a direct effect of temperature on the binding efficiencies of the membrane pumps similar to the temperature-induced  $K_m$  changes of enzymes in fish and plants (Somero & Low, 1976; Teeri & Peet, 1978). The decline in pump efficiency at either the plasma membrane or tonoplast membrane could result in an alteration of ion transport activity resulting in an elevation of salts (i.e. calcium) and metals (e.g. aluminium, iron, copper and cadmium) in the cytoplasm. Most, if not all, of the other stresses reported to induce the heat-shock response would also affect either the gradient of metals, the energy required to drive the membrane pumps or the transport of salts across the membrane. The leakage of calcium across the membrane would result in the activation of calmodulin [CAM] (Fig. 1, [1]) and, ultimately, enzyme activities associated with

calmodulin binding (i.e. membrane ATPases (Fig. 1, [2]), kinases, and phospholipase D (Fig. 1, [3])). The potential activation of phospholipase D could account for the loss of the endoplasmic reticulum's phospholipid membrane during the initial phases of heat-shock (Fig. 1, [4]) (Belanger *et al.*, 1986). The activity of the phospholipase would decline as the calmodulin-activated membrane ATPases removed excess ions from the cytoplasm. Calcium influx into corn roots has previously been reported as a result of cold shock (Zocchi & Hanson, 1982). Zocchi & Hanson (1983) suggested that a  $\text{Ca}^{2+}/\text{H}^{+}$  exchange functions normally to maintain very low  $\text{Ca}^{2+}$  concentrations, with a  $\text{Ca}^{2+}$ -ATPase activated only when  $\text{Ca}^{2+}$  levels rise and calmodulin is activated. A similar activation may occur in response to the heat-shock temperatures.

Leakage of metals into cells can be extremely toxic. Cells have evolved mechanisms to aid in scavenging intracellular metals (Rausser, 1981). Glutathione functions as a cellular reductant and has been shown to be involved in removal of cellular toxins (herbicides, metals and so on) (Cherian & Goyer, 1978; Grill, Winnacker & Zenk, 1986). If declining energy levels or direct temperature effects on the kinetic constants of membrane pumps alter the intracellular concentrations of calcium and metals, then increased glutathione concentrations could aid in cellular detoxification (Fig. 1, [5]). To overcome high-temperature-induced kinetic changes in the enzyme responsible for maintaining glutathione in the reduced state (*viz.* glutathione reductase) (Mahan, Burke & Orzech, 1987), elevated levels of glutathione would be required for the maintenance of glutathione reductase activity. The increase in glutathione concentrations reported in high temperature stressed maize (Nieto-Sotelo & Ho, 1986) is consistent with this model. The mechanism responsible for the thermally induced increase in glutathione content has not been identified, however, it may require the synthesis of the enzymes  $\gamma$ -glutamylcysteine synthetase and glutathione synthase. Some metals have been shown to induce the synthesis of linear polymers of glutathione, termed phytochelatins (Grill, Winnacker & Zenk, 1985). These phytochelatins function as metal-sequestering peptides via metal-thiolate co-ordination and have been shown to be induced by a range of metals (Grill *et al.*, 1986). Although specific metal transporters have not been isolated, permeability studies by Gutknecht (1983) suggest that  $\text{Cd}^{2+}$  transport across membranes is protein mediated and correlates with  $\text{Cd}^{2+}$ , not  $\text{CdCl}_2$ , concentration.

Concomitant with the initial scavenging of the metals by glutathione and phytochelatins, some of the metals may interact with genes containing similar sequences to the metal ion responsive elements found upstream in metallothionein genes of animals (Fig. 1, [6]) (Serfling *et al.*, 1985). Similar sequences have been reported far upstream in the Gmhsp 17.5-E




Figure 1. Model of potential cellular responses to high temperature stress. Key reactions [1 to 8] are discussed in the text.

(*Glycine max* heat-shock protein, 17.5 kDa) genes (Czarnecka *et al.*, 1985). The degree to which the metals would be available for HSP induction would be related to the level of glutathione, cysteine, and phytochelatins in the cell. This may explain in part the observed changes in the induction temperature of the heat-shock response following several cycles of heat-shock (Key *et al.*, 1985). The reported increases in glutathione content in response to the high temperature stress would provide increased protection from the metals, thereby functionally removing them from HSP induction.

The mechanism regulating the changes in the specificity of ribosomes for the heat-shock mRNA is unknown. The association of a 22 kD (P22) cytoplasmic protein has been reported during the transition to selective mRNA translation (McMullin & Hallberg, 1986). The mechanism of interaction between the P22 and the ribosome, however, remains unknown. It is possible that the binding of the P22 to free ribosomes is associated with metal binding during the thermally-induced leakage of metals into the cell or binding may be related to protein phosphorylation by a calmodulin-activated kinase (Fig. 1, [7]).

Finally, some of the small molecular weight HSP may function in chelation of metals (Fig. 1, [8]). Most metal-chelating proteins that have been identified are low molecular weight proteins whose synthesis is induced by specific metals (Weigel & Jager, 1980). The low-molecular-weight heat-shock proteins have also been shown to be induced by metals, with some specificity between the metal tested and the appearance of mRNA for particular low molecular weight HSP (Czarnecka *et al.*, 1984). These chelating proteins could be protected from proteolysis while binding the metal and become susceptible to proteolysis following removal of the metal. This hypothesis is consistent with the observation that HSP do not accumulate to significant levels following a high-temperature treatment, yet have been shown to accumulate in field-stressed plants where elevation in leaf temperatures is associated with tissue dehydration and water stress (Burke *et al.*, 1985). Water stress can result in tissue dehydration which would increase the cellular ion concentrations, thereby requiring their functional removal via chelation by low molecular weight HSP until the ions could be transported out of the cell or until the stress was alleviated.

In summary, the present authors have presented a working model of possible responses of higher-plant cells to elevated temperatures. Because of the reported induction of heat-shock proteins by metals, and because of the known sensitivity of enzyme systems to temperature changes, the reported cellular responses presented in this model cannot be dismissed based upon the existing heat-shock response literature. Future research on heat-shock

responses of higher plants should consider the possibilities addressed in this research model.

## References

- Barnett, T., Altschuler, M., Daniel, C.N. & Mascarenhas, J.P. (1980) Heat shock induced proteins in plant cells. *Developmental Genetics* **1**, 331–340.
- Baszczynski, C.L., Walden, D.B. & Atkinson, B.G. (1983) Regulation of gene expression in corn (*Zea mays* L.) by heat shock. II: *In vitro* analysis of RNAs from heat-shocked seedlings. *Canadian Journal of Biochemistry and Cell Biology*, **61**, 395–403.
- Baszczynski, C.L., Walden, D.B. & Atkinson, B.G. (1985) Maize genome response to thermal shifts. In *Changes in Eukaryotic Gene Expression in Response to Environmental Stress* (eds B. G. Atkinson & D. B. Walden), pp. 349–371. Academic Press Inc, New York.
- Belanger, F.C., Brodl, M.R. & Ho, T.H.-D. (1986) Heat shock causes destabilization of specific mRNAs and destruction of endoplasmic reticulum in barley aleurone cells. *Proceedings of the National Academy of Sciences (USA)*, **83**, 1354–1358.
- Bewley, D.J., Larsen, K.M. & Papp, J.T. (1983) Water stress induced changes in the pattern of protein synthesis in maize seedling mesocotyls: a comparison with the effects of heat shock. *Journal of Experimental Botany*, **34**, 1126–1133.
- Bienz, M. (1984) Developmental control of the heat shock response in *Xenopus*. *Proceedings of the National Academy of Sciences*, **81**, 3138.
- Burke, J.J., Hatfield, J.L., Klein, R.R. & Mullet, J.E. (1985) Accumulation of heat shock proteins in field-grown cotton. *Plant Physiology*, **78**, 394–398.
- Chen, Y.-M., Kamisaka, S. & Masuda, Y. (1986) Enhancing effects of heat shock and gibberellic acid on the thermotolerance in etiolated *Vigna radiata*. I: Physiological aspects of thermotolerance. *Physiologia Plantarum*, **66**, 595–601.
- Cherian, M.G. & Goyer, R.A. (1978) Metallothioneins and their role in metabolism and toxicity of metals. *Life Science*, **23**, 1–10.
- Cooper, P. & Ho, T.H.-D. (1983) Heat shock proteins in maize. *Plant Physiology* **71**, 215–222.
- Cooper, P. & Ho, T.H.-D. (1984) The specificity of the heat shock response in maize. *Plant Physiology* **75**, 432–441.
- Cooper, P. & Ho, T.H.-D. (1987) Intracellular localization of heat shock proteins in maize. *Plant Physiology*, **84**, 1197–1203.
- Craig, E.A. (1986) The heat shock response. *CRC Critical Reviews in Biochemistry*, **18**, 239–280.
- Czarnecka, E., Edelman, L., Schoffl, F. & Key, J.L. (1984) Comparative analysis of physical stress responses in soybean seedlings using cloned heat shock cDNAs. *Plant Molecular Biology*, **3**, 45–58.
- Czarnecka, E., Gurley, W.B., Nagao, R.T., Mosquera, L.A. & Key, J.L. (1985) DNA sequence and transcript mapping of a soybean gene encoding a small heat shock protein. *Proceedings of the National Academy of Sciences (USA)*, **82**, 3726–3730.
- Ginzburg, C. & Salomon, R. (1986) The Effect of Dormancy on the Heat shock response in *Gladiolus* Cormels. *Plant Physiology*, **81**, 259–267.
- Grill, E., Winnacker, E.-L. & Zenk, M.H. (1985) Phytochelatins: the principal heavy-metal complexing peptides of higher plants. *Science*, **230**, 674–676.
- Grill, E., Winnacker, E.-L. & Zenk, M.H. (1986) Synthesis of seven different homologous phytochelatins in metal-exposed *Schizosaccharomyces pombe* cells. *FEBS Letters*, **197**, 115–120.
- Gutknecht, J. (1983) Cadmium and thallous ion permeabilities through lipid bilayer membranes. *Biochimica et Biophysica Acta*, **735**, 185–188.
- Hadwiger, L.A. & Wagoner, W. (1983) Effect of heat shock on the mRNA-directed disease resistance response of peas. *Plant Physiology*, **72**, 553–556.

- Heikkilä, J.J., Papp, J.E.T., Schultz, G.A. & Bewley, J.D. (1984) Induction of heat shock protein messenger RNA in maize mesocotyls by water stress, abscisic acid, and wounding. *Plant Physiology*, **76**, 270–274.
- Hong-Qi, Z., Croes, A.F. & Linskens, H.F. (1984) Qualitative changes in protein synthesis in germinating pollen of *Lilium longiflorum* after a heat shock. *Plant, Cell and Environment*, **7**, 689–691.
- Key, J.L., Czarnecka, E., Lin, C.Y., Kimpel, J., Mothershed, C. & Schoffl, F. (1983) A comparative analysis of the heat shock response in crop plants. In *Current Topics in Plant Biochemistry and Physiology*, Vol. 2. (eds D. D. Randall, D. G. Blevins, R. L. Larson & B. J. Rapp), pp. 107–118. University of Missouri, Columbia, Missouri.
- Key, J. L., Kimpel, J.A., Vierling, E., Lin, C.Y., Nagao, R.T. & Czarnecka, E. (1985) Physiological and molecular analyses of the heat shock response in plants. In *Changes in Eukaryotic Gene Expression in Response to Environmental Stress* (eds B. G. Atkinson & D. B. Walden), pp. 327–348. Academic Press, New York.
- Key, J.L., Lin, C.Y., Ceglaz, E. & Schoffl, F. (1982) The heat shock response in plants. In *Heat Shock, from Bacteria to Man*. (eds M. J. Schlesinger, M. Ashburner & A. Tissieres), p. 329. Cold Spring Harbor Lab., Cold Spring Harbor, New York.
- Key, J. L., Lin, C.Y. & Chen, Y.M. (1981) Heat shock proteins of higher plants. *Proceedings of the National Academy of Sciences (USA)*, **78**, 3526–3530.
- Lim, L., Hall, C., Leung, T. & Whatley, S. (1984) The relationship of the rat brain 68 kDa microtubule-associated protein with synaptosomal plasma membranes and with the *Drosophila* 70 kDa heat-shock protein. *Biochemical Journal*, **224**, 677–680.
- Lindquist, S. (1986) The heat-shock response. *Annual Review of Biochemistry* 1986, **55**, 1151–1191.
- Mahan, J.R., Burke, J.J. & Orzech, K.A. (1987) The 'thermal kinetic window' as an indicator of optimum plant temperature. *Plant Physiology*, **83**, S–87.
- Mansfield, M.A. & Key, J.L. (1987) Synthesis of low molecular weight heat shock proteins in plants. *Plant Physiology*, **84**, 1007–1017.
- McMullin, T.W. & Hallberg, R.L. (1986) Effect of heat shock on ribosome structure: appearance of a new ribosome-associated protein. *Molecular and Cellular Biology*, **6**, 2527–2535.
- Meyer, Y. & Chartier, Y. (1983) Long-lived and short-lived heat-shock proteins in tobacco mesophyll protoplasts. *Plant Physiology*, **72**, 26–32.
- Nieto-Sotelo, J. & Ho, T.H.-D. (1986) Effect of heat shock on the metabolism of glutathione in maize roots. *Plant Physiology*, **82**, 1031–1035.
- Nover, L. (ed.) (1984) *Heat Shock Response of Eukaryotic Cells*. VEB Georg Thieme, Leipzig.
- Nover, L. & Scharf, K.D. (1984) Synthesis, modification and structural binding of heat-shock proteins in tomato cell cultures. *European Journal of Biochemistry*, **139**, 303–313.
- Nover, L., Scharf, K.D. & Neumann, D. (1983) Formation of cytoplasmic heat shock granules in tomato cell cultures and leaves. *Molecular and Cellular Biology*, **Sept**, 1648–1655.
- Ougham, H.J. & Stoddart, J.L. (1986) Synthesis of heat-shock protein and acquisition of thermotolerance in high-temperature tolerant and high-temperature susceptible lines of sorghum. *Plant Science*, **44**, 163–167.
- Pitto, L., Schiavo, F.L., Giuliano, G. & Terzi, M. (1983) Analysis of the heat-shock protein pattern during somatic embryogenesis of carrot. *Plant Molecular Biology*, **2**, 231–237.
- Rausser, W.E. (1981) Occurrence of metal binding proteins in plants. In *International Conference Heavy Metals in the Environment*, pp. 281–284. CEP Consultants, Edinburgh.
- Schoffl, F. & Baumann, G. (1985) Thermo-induced transcripts of a soybean heat shock gene after transfer into sunflower using a Ti plasmid vector. *EMBO Journal*, **4**, 1119–1124.
- Serfling, E., Lubbe, A., Dorsch-Hasler, K. & Schaffner, W. (1985) Metal-dependent SV40 viruses containing inducible enhancers from the upstream region of metallothionein genes. *EMBO Journal*, **4**, 3851–3859.
- Sinibaldi, R.M. & Turpen, T. (1985) A heat shock protein is encoded within mitochondria of higher plants. *Journal of Biological Chemistry*, **260**, 15382–15385.
- Somero, G.N. & Low, P.S. (1976) Temperature: a 'shaping force' in protein evolution. *Biochemical Society Symposium*, **41**, 33–42.
- Teeri, J.A. & Peet, M.M. (1978) Adaptation of malate dehydrogenase to environmental temperature variability in two populations of *Potentilla glandulosa* Lindl. *Oecologia (Berlin)*, **34**, 133–141.
- Vierling, E., Mishkind, M.L., Schmidt, G.W. & Key, J.L. (1986) Specific heat shock proteins and transport into chloroplasts. *PNAS (USA)*, **83**, 361–365.
- Vincent, M. & Tanguay, R.M. (1979) Heat-shock induced proteins present in the cell nucleus of chironomus tentans salivary gland. *Nature*, **281**, 501–503.
- Weigel, H.J. & Jager, H.J. (1980) Subcellular distribution and chemical form of cadmium in bean plants. *Plant Physiology*, **65**, 480–482.
- Xiao, C.-M. & Mascarenhas, J.P. (1985) High temperature-induced thermotolerance in pollen tubes of *Tradescantia* and heat-shock Proteins. *Plant Physiology*, **78**, 887–890.
- Zocchi, G. & Hanson, J.B. (1982) Calcium influx into corn roots as a result of cold shock. *Plant Physiology*, **70**, 318–319.
- Zocchi, G. & Hanson, J.B. (1983) Calcium transport and ATPase activity in a microsomal vesicle fraction from corn roots. *Plant, Cell and Environment*, **6**, 202–209.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.